

10 REFERENCES

- Arooz, R.H., and Arshad, M.A., 1996; Soil infiltration and hydraulic conductivity under long-term no-tillage and conventional tillage systems. *Canadian Journal of Soil Science*. 76: 143-152
- Tiessen, K.H.D., Elliott, J.A., Yarotski, J., Lobb, D.A., Flaten, D.N., and Glozier, N.E. , 2010. Conventional and Conservation Tillage: Influence on Seasonal Runoff, Sediment, and Nutrient Losses in the Canadian Prairies. *J. Environ. Qual.* 39:964–980.
- Barry-Eaton District Health Department. 2011. Time of Sale or Transfer (TOST) Program: The first three years. <http://www.barryeatonhealth.org/Portals/9/EH/FIRST%20THREE%20YEARS%20OF%20TOST.pdf> . Accessed November 30, 2016.
- Beletsky, D., Saylor, J.H., and Schwab, D.J. 1999. Mean Circulation in the Great Lakes. *J. Great Lakes Res.* 25 (1): 78-93.
- Edly, K. and Wuycheck, J. 2006. Water Quality and Pollution Control in Michigan: Sections 303(d), 305(b), and 314 Integrated Report. MDEQ Report #MI/DEQ/WB-06/019.
- Garret, E. 2016. Michigan adds algae-choked Lake Erie to list of 'impaired waters'. www.mlive.com/news/index.ssf/2016/11/michigan_lake_erie_impaired_ep.html
- Global Climate Change Impacts in the United States, Thomas R. Karl, Jerry M. Melillo, and Thomas C. Peterson,(eds.). Cambridge University Press, 2009.
- Hooper, Frank F. and S. L. Kohler. 2000. Measurement of stream velocity and discharge. Chapter 19 in Schneider, James C. (ed.) 2000. *Manual of fisheries survey methods II: with periodic updates*. Michigan Department of Natural Resources, Fisheries Special Report 25, Ann Arbor.
- International Joint Commission (2014). *A Balanced Diet for Lake Erie: Reducing Phosphorus Loadings and Harmful Algal Blooms*. Report of the Lake Erie Ecosystem Priority.
- Lick, W., Lick, J., and Ziegler, C.K. 1994. The Resuspension and Transport of Fine-Grained Sediments in Lake Erie. *J. Great Lakes Res.* 20 (4): 599-612
- Liu, et al. 2006. Modeling the Transport and Inactivation of *E.coli* and Enterococci in the Near-Shore Region of Lake Michigan. *Environ, Sci. Technol.* 40: 5022-5028.
- "Long Term Hydrologic Impact Analysis (L-thia)." *Engineering.purdue.com*. Purdue University. Web. 9 Mar. 2016. <<https://engineering.purdue.edu/~lthia/>>.
- Michalak, A.M., et al. 2013. Record-setting algal bloom in Lake Erie caused by agricultural and meteorological trends consistent with expected future conditions. *PNAS* 110 (16): 6448-6452.

Michigan Department of Environmental Quality. 2017. State of Michigan Domestic Action Plan for Lake Erie. [http://www.michigan.gov/documents/deq/Michigan Domestic Action Plan](http://www.michigan.gov/documents/deq/Michigan_Domestic_Action_Plan).

Michigan Department of Environmental Quality. 2014. Lapointe Drain Watershed Landscape Level Wetland Functional Assessment (Enhanced NWI).

Michigan Office of the Great Lakes, 2016. Michigan State of the Great Lakes.

Mozola, A.J. 1970. Geology for environmental planning in Monroe County, Michigan: Michigan Geological Survey Report of Investigation 13, 34

Murray, et al. 2011. Canine Scent and Microbial Source Tracking in Santa Barbara, California. WERF.

Omerink, J. and A. Gallant, 1988. Ecoregions of the Upper Midwest States. USEPA Environmental Research Laboratory. EPA/600/3-88/037

Ripke, M. 2007. Total Maximum Daily Load for *E. coli* for Lake Erie Luna Pier Beach, Monroe County. Surface Water Assessment Section, Water Bureau, Michigan Department of Environmental Quality.

Schmitt, G. and Sunday, E. 2007. Total Maximum Daily Load for Dissolved Oxygen and Total Phosphorus for S.S. Lapointe Drain, Monroe County. Surface Water Assessment Section, Water Bureau, Michigan Department of Environmental Quality.

Schottler, S.P., et al. 2013. Twentieth century agricultural drainage creates more erosive rivers. Hydrological Processes 28 (4): 1951-1961.

Smith, R.A., R.B. Alexander, G.E. Schwarz. 2003. Natural Background Concentrations of Nutrients in Streams and Rivers of the Conterminous United States. Environmental Science and Technology 37(14): 3039-3047.

STEPL Input Data Server Version 1.0. Web. 9 Mar. 2016. <<http://it.tetratex.com/steplweb/steplweb.html>>. LaPlaisance Creek-Frontal Lake Erie Watershed

Tetra Tech Inc. User's Guide Spreadsheet Tool for the Estimation of Pollutant Load (STEPL). 2011 ed. Fairfax, VA: Tetra Tech, 2011. U.S. Environmental Protection Agency, Aug. 2011. Web.

Turner, C. and Bobrin, J., 2006. The Application of Source Tracking Methods in a Municipal Storm Sewer System to support TMDL Implementation. Presentation at StormCon, Denver, Colorado.

United States Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS), 2014. Regional Conservation Partnership Program - Great Lakes Region Introduction. <https://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/programs/farmbill/rcpp/?cid=stelprdb1254125>

United States Environmental Protection Agency (USEPA). Wetlands Factsheet.

United States Environmental Protection Agency (USEPA). 1997. Volunteer Stream Monitoring: A Methods Manual

Verhougstraete, et.al., 2015. Linking fecal bacteria in rivers to landscape, geochemical, and hydrologic factors and sources at the basin scale. PNAS 112 (33): 10419-10424

Wilson, J.L., Schumacher, J.G., and Burken, J.G. 2014. Occurrence and origin of Escherichia coli in water and sediment at two public swimming beaches at Lake of the Ozarks State Park, Camden County, Missouri, 2011-13: U.S. Geological Survey Scientific Investigations Report 2014-5005, 59p.
<http://dx.doi.org/10.3133/sir20145005>